

A TAXONOMIC REVISION OF *OCTOPUS AUSTRALIS* HOYLE, 1885
(OCTOPODIDAE: CEPHALOPODA), WITH A REDESCRIPTION OF
THE SPECIES

By R. W. TAIT

Department of Invertebrate Zoology, National Museum of Victoria,
285-321 Russell Street, Melbourne, Victoria, 3000

Abstract

Octopus australis Hoyle from South Eastern Australian waters is fully redescribed. Several species from New Zealand previously synonymised with it are recognised as distinct and their nomenclatural status is discussed.

Introduction

Octopus australis was described by Hoyle, based on one female and one immature specimen from Port Jackson, N.S.W. Subsequently, Massey (1916), Robson (1929), Benham (1942) and Dell (1952) have described material from New Zealand, which they have synonymised with *O. australis* Hoyle. Dell's synonymy includes *Polypus campbelli* Smith, 1902; *Polypus australis* Massey, 1916; *Polypus* cf *australis* Berry, 1918; and *Robsonella australis* Benham, 1942. Specimens from South Eastern Australian waters match closely with the brief type description, but differ to the descriptions of Smith, Massey, Robson, Benham and Dell. To clarify this situation, the type specimens of *O. australis* were borrowed from the British Museum for comparison with other Australian and New Zealand material.

The identity of the S.E. Australian species is confirmed as *Octopus australis* Hoyle, and a complete redescription is given. The New Zealand species *Polypus campbelli* Smith, *Polypus australis* Massey and *Robsonella*

australis Benham are identified as a separate species group, and their nomenclatural status is discussed.

Measurements and Abbreviations

The measurements and abbreviations used are the same as given in Voss (1963), with the exception of head length. Head length (H.L.) is taken from the junction of the dorsal pair of arms to the midpoint between the eyes. Measurements are given in Table 1, indices are expressed in Table 2. Other abbreviations used are BM(NH)—British Museum (Natural History); NMV—National Museum of Victoria.

***Octopus australis* Hoyle, 1885**

Plate 1—a, b Figures 1-2.

1885a *Octopus australis* Hoyle, p. 224.

1885b *Octopus australis* Hoyle, pp. 98-99.

1886 *Octopus australis* Hoyle, pp. 88-89, pl. 3, figs. 4-5.

Materials examined:

Sex	M.L. (mm)	Reg. No.	Location	Date coll.	Depth (m)
Holotype ♀	22	BM(NH)1889.4.24.28.9	Port Jackson, N.S.W.	April 1874	11-28
Paratype ♂	12	BM(NH)1889.4.24.28.9	Port Jackson, N.S.W.	April 1874	11-28
Other material					
♂	16	NMV F25247	Western Port Bay, Vic.	—	—
♂	21	NMV F30860	40°34'S, 144°46'E	4. 2.1981	68
♂	27	NMV F31265	37°55'S, 144°58'E	18. 3.1980	7
♂	37	NMV F31003	32°24'S, 133°30'E	23. 8.1973	49
♂	42	NMV F31265	37°55'S, 144°58'E	18. 3.1980	7

Sex	M.L. (mm)	Reg. No.	Location	Date coll.	Depth (m)
<i>Materials</i>					
♂	45	NMV F21911	38°02'S, 145°04'E	1961	11
♂	45	NMV F21911	38°02'S, 145°04'E	1961	11
♂	46	NMV F31267	32°24'S, 133°24'E	26.10.1973	40
♂	54	NMV F31002	39°38'S, 145°06'E	3. 2.1981	66
♂	56	NMV F21911	38°02'S, 145°04'E	1961	11
♂	67	NMV F31260	32°13'S, 133°52'E	27. 4.1973	8
♂	67	NMV F25436	38°07'S, 145°06'E	1964	—
♂	72	NMV F31002	39°38'S, 145°06'E	3. 2.1981	66
♂	73	NMV F31264	38°03'S, 145°06'E	7. 6.1978	—
♀	9	NMV F31263	38°55'S, 145°55'E	9.11.1972	12
♀	14	NMV F31262	35°23'S, 137°17'E	21. 1.1971	54
♀	17	NMV F25247	Western Port Bay, Vic.	—	—
♀	17	NMV F30927	40°50'S, 146°07'E	4. 2.1981	66
♀	19	NMV F25247	Western Port Bay, Vic.	—	—
♀	25	NMV F31265	35°55'S, 144°58'E	18. 3.1980	7
♀	28	NMV F31265	37°55'S, 144°58'E	18. 3.1980	7
♀	34	NMV F31265	37°55'S, 144°58'E	18. 3.1980	7
♀	34	NMV F31003	32°24'S, 133°30'E	23. 8.1973	49
♀	37	NMV F31265	37°55'S, 144°58'E	18. 3.1980	7
♀	37	NMV F31003	32°24'S, 133°30'E	23. 8.1973	49
♀	41	NMV F24485	37°51'S, 144°57'E	15. 1.1930	—
♀	44	NMV F25245	Western Port Bay, Vic.	1963	—
♀	47	NMV F24492	37°59'S, 145°01'E	10.11.1973	—
♀	49	NMV F31002	39°38'S, 145°06'E	3. 2.1981	66
♀	49	NMV F31002	39°38'S, 145°06'E	3. 2.1981	66
♀	62	NMV F31265	37°55'S, 144°58'E	18. 3.1980	7
♀	88	NMV F24437	38°13'S, 145°02'E	3.10.1957	—

Diagnosis

Size up to 90 mm M.L., arms long, mantle sculpture fine dorsally, smooth ventrally, lateral integumental ridge usually present. 7-8 gill lamellae in outer demibranch; funnel organ with closely opposed, occasionally partially fused VV units. Hectocotylied arm 58-75% of A_{LIII} length; ligula robust, 12-18% of arm length, with double row of fine papillae along median oral excavation.

Description

Mantle globular, quite broad, well demarked from head; mantle aperture wide; head narrow; eyes small, protuberant (Fig. 1a, Plate 1b). Funnel free for about half its length; funnel organ with two closely opposed V shaped units, ventral and dorsal limbs of approximately equal lengths; units may be partially fused medially (Figs. 1, b-d).


Arms long, subequal, tapering to fine tips; suckers moderate in size, no enlarged suckers in males. Web shallow, extends up the ventral side of the arms for almost their entire length. Web formula D.C.B.E.A. to C.D.B.A.E., dorsal and ventral sectors always shallowest.

Third right arm of males hectocotylied, 58-75% length of its opposite member; spermatophoral groove well developed but without any conspicuous thickening of the interbrachial web. Ligula large (Figs, 1e, f), deeply excavated, usually curved orally; medially two rows of very small papillae are present along the excavation. Calamus short, acutely pointed.


Gills moderate in length, outer demibranch with 7-8 primary lamellae, plus a terminal lamella.

Reproductive system of males typical of the genus; penis (Fig. 2a) long with a single coiled diverticulum on the right hand side, genital aperture subterminal, on right hand side. Spermatophores (Fig. 2, b-e) long, thin; horn with 2-3 coils close to oral end, oral cap expanded. Female reproductive system without distinctive features; eggs large, length 9-14 mm, attached singly by a stalk approximately 8 mm in length; each clutch of 80-130 eggs (Tait 1980).


Fig. 1. *Octopus australis* Hoyle, 1885, a. Ventral view, male, NMV F31267, 46 mm M.L. b-d. Funnel organs. b. Holotype, female, 22 mm M.L. c. NMV F31265, female, 28 mm M.L. d. NMV F31265, male, 42 mm M.L. e-f. Ligula. NMV F31267, 46 mm M.L. g. Radula. Holotype.




a



b




c



d




e



f



g



Alimentary canal of normal octopodan type; crop with an anterior caecum of about 10% of its length; posterior oesophagus short; posterior salivary glands elongate, connect to buccal mass by a common duct, ducts to crop separate. Stomach bipartite; caecum strongly coiled; ducts to stomach and caecum originate separately from hepatopancreas, intestine without conspicuous differentiation. Ink sac large, embedded in surface of hepatopancreas, connected to intestine near anus by a short duct.

Beaks (Fig. 2f) typically octopodan; dorsal rostrum curved, wings transparent in small individuals; ventral beak with very blunt rostrum, wings with small anterior protuberances.

Radula (Fig. 1g) with B₃₋₄ seriation (holotype B₄); rhachidian tooth asymmetrical, 1-2 cusps on each side; first laterals with one sharp cusp, second laterals with one cusp, third laterals long, straight or slightly curved; marginal plates oblong, elongate.

Dorsal mantle surface covered by fine tubercles, supraocular cirri often present, ventral surface smooth. Depending on condition of preservation, a ventro-lateral integumental ridge may be present (Fig. 1a, Plate b), most evident adjacent to mantle aperture, often disappearing posteriorly. In live animals this may be extended into a shallow web, or evident only as a fine, light coloured line.

Colour of preserved specimens brown to purplish dorsally, ventral surface cream. Faint pair of roughly circular ocelli present in some specimens posterior to the eyes; each ocellus comprises a dark ring with a lighter centre; they are most apparent in live animals and fade during preservation.

Colour of preserved specimens brown to purplish dorsally, ventral surface cream. Faint pair of roughly circular ocelli present in some specimens posterior to the eyes; each ocellus comprises a dark ring with a lighter centre; they are most apparent in live animals and fade during preservation.

Males have spermatophores in the Needhams Sac when larger than 20-25 mm M.L. (10-13 g), and may reach 70-80 mm M.L. (250 g). Females have large, white and translucent eggs in the ovary when larger than 30-40 mm M.L. (40-60 g), and do not usually grow beyond 50 mm M.L. (100 g).

Fig. 2. *Octopus australis* Hoyle, 1885. a. Penis. NMV F31002, 72 mm M.L. b-e. Spermatophore. NMV F31002, 54 mm M.L. b. Whole spermatophore. c. Cement body. d. Oral end of cement body. e. Oral end. f. Beaks. NMV F31002, female, 49 mm M.L.

Distribution

The collections of the National Museum of Victoria contain specimens of this species from New South Wales, Victorian, Tasmanian and South Australian waters, to depths of 70 m.

Discussion

The confusion relating to the identity of *Octopus australis* Hoyle is due largely to the lack of a mature male type specimen. Although their external morphology is somewhat similar, New Zealand and Australian species may be readily separated by the form of the hectocotylus. Details of the funnel organ, radula, surface sculpture and number of gill lamellae of the holotype indicate that it is conspecific with the Australian material studied.

Of the four species synonymised with *Octopus australis* Hoyle in Dell (1952), detailed descriptions exist for three. Robson (1929) redescribed the holotype of *Polypus campbelli* Smith, from Campbell Island (N.Z.) as having a W-type funnel organ, 10 lamellae in each gill demibranch, enlarged suckers in the male and a L.L.I. of only 8.5%. Massey's (1916) *Polypus australis*, from New Zealand, has a L.L.I. of 11%, a W type funnel organ and symmetrical seriation of the radula. *Robsonella australis*, also from New Zealand, was described by Benham (1942) and Dell (1952). It has a W type funnel organ, stronger cusps on the radula than *O. australis* Hoyle, and eggs of only 2.5-2.8 mm in length (Brough 1965). Further, I have remeasured the five mature males described by Benham and one other from 38°10'S, 147°49'E (NMV F31259) and these have the hectocotylus indices given in Table 3.

TABLE 3

Hectocotylus indices of *Robsonella australis* Benham

	n	mean	S.D.(n-1)	range
HcAI	5*	77.5	3.5	73-81
LLI	6	8.1	1.5	6-10
CLI	6	39.1	7.4	33-53

* A_{LI} of one specimen regenerating.

These are quite distinct from the corresponding indices given for *O. australis* Hoyle in Table 2. Therefore, all the New Zealand species previously considered to be synonyms of *O. australis* Hoyle appear to be separate and distinct. The description of *Polypus* cf. *australis* from South East Australia, by Berry (1918), is not detailed enough to confirm his tentative identification.

Robson (1929) synonymised *Polypus campbelli* Smith and *Octopus australis* Hoyle by comparing, in part, the ligula of each. Robson's material included the types of *O. australis* Hoyle and Massey's *Polypus australis*. As a mature male type is lacking, Robson's synonymy was therefore based on Massey's material, already shown to be distinct from *O. australis* Hoyle. Furthermore, Robson's description of the radula of the *O. australis* holotype could not have been from the types, as both type specimens had buccal masses intact when loaned by the British Museum.

Nomenclatural Status of the New Zealand Species

The New Zealand species *Polypus australis* Massey and *Robsonella australis* Benham are both junior homonyms of *Octopus australis* Hoyle. This follows the renaming of the genus *Polypus* as *Octopus* by Robson (1929), and the regrouping of *Robsonella* under *Octopus* by Pickford (1955). If the synonymy of these two species and *Polypus campbelli* Smith, as given in Benham (1942) and Dell (1952) is correct, then *Octopus campbelli* (Smith) is the correct senior synonym. However, the differences in the radula of *P. australis* Massey and the enlarged suckers in male *P. campbelli* Smith make it probable that these species and *R. australis* Benham are distinct from each other. *Robsonella australis* Benham and *Polypus australis* Massey must then be renamed to prevent the occurrence of two homonyms of *Octopus australis* in close geographic proximity. A review of this New Zealand species group is urgently required to remove this problem.

Acknowledgements

I wish to thank Dr. C. C. Lu, Curator,

Department of Invertebrate Zoology, National Museum of Victoria, for his advice and reading of the manuscript. For providing material either by loan or donation I gratefully acknowledge: Mr. F. Naggs, British Museum (Natural History); Dr. A. C. Harris, Otago Museum, New Zealand and the National Museum of Victoria Bass Strait Survey. Funding for the Bass Strait Survey was generously provided by A.M.S.T.A.C. Thanks also to Mrs. J. Andrewartha who provided the illustrations and Mrs. L. Anderson for typing the manuscript.

References

- BENHAM, W. B., 1942. The octopodous Mollusca of New Zealand. 1. The midget octopus of the coastal waters. *Trans. R. Soc. N.Z.* 72: 226-236.
- BERRY, S. S., 1918. Report on the Cephalopoda obtained by the F.I.S. 'Endeavour' in the Great Australian Bight and other Southern Australian localities. *Zool. (biol.) Results Fish. Exp. 'Endeavour'*. 4(5): 203-298, pls. 59-88.
- BROUGH, E. J., 1965. Egg care, eggs and larvae in the midget octopus *Robsonella australis* (Hoyle). *Trans. R. Soc. N.Z. Zool.*, 6: 7-19.
- DELL, R. K., 1952. The recent Cephalopoda of New Zealand. *Dom. Mus. Bull.* 16: 157 pp., 35 pls.
- HOYLE, W. E., 1885a. Diagnoses of new species of Cephalopoda collected during the cruise of H.M.S. 'Challenger'. Part 1. The Octopoda. *Ann. Mag. nat. Hist.*, ser. 5, 15: 222-236.
- HOYLE, W. E., 1885b. Preliminary report on the Cephalopoda collected during the cruise of H.M.S. 'Challenger'. Part I. The Octopoda. *Proc. R. Soc. Edinb.* 13: 94-114.
- HOYLE, W. E., 1886. Report on the Cephalopoda collected by H.M.S. 'Challenger' during the years 1873-76. *Rep. scient. Results Voy. Challenger Zool.* 16(44): 245 pp., 33 pls.
- MASSEY, A. L., 1916. Cephalopoda. *Br. Antarct. Terra Nova Exped. 1910 Zool.* 2(7): 141-175.
- PICKFORD, G. E., 1955. A revision of the Octopodinae in the collections of the British Museum. *Bull. Br. Mus. nat. Hist. Zool.* 3(3): 151-167.
- ROBSON, G. E., 1929. *A monograph of the recent Cephalopoda, based on the collections in the British Museum (Natural History). Part I, Octopodinae.* London.
- TAIT, R. W. 1980. *Aspects of the ecology and life history of Octopus australis Hoyle, from Northern Port Phillip Bay.* B.Sc.(Hons) thesis, Monash University, Clayton, 69 pp., 13 pls.
- VOSS, G. L., 1963. Cephalopods of the Philippine Islands. *Bull. U.S. natn. Mus.* 234: 180 pp.

TABLE 1
Measurements (in mm) of *Octopus australis* Hoyle

BM(NH) 1889.4.24.28.9	NMV F25436	NMV F31265
NMV F25247	NMV F31002	NMV F31003
NMV F30860	NMV F31264	NMV F24485
NMV F31265	NMV F31263	NMV F25245
NMV F31003	NMV F31262	NMV F24492
NMV F31265	NMV F25247	NMV F31002
NMV F21911	NMV F30927	NMV F31002
NMV F21911	NMV F25247	NMV F31265
NMV F31267	BM(NH) 1889.4.24.28.9	NMV F24437
NMV F31002	NMV F31265	
NMV F21911	NMV F31265	
NMV F31260	NMV F31265	
	NMV F31003	

Sex	♂	♂	♂	♂	♂	♂	♂	♂	♂	♂	♂	♂
M.L.	12	16	21	27	37	42	45	45	46	54	56	67
Tot. L.	38	47	86	107	170	184	158	163	220	280	200	245
MW	10	12	17	21	29	31	33	27	35	48	35	36
H.L.	3	5	9	10	17	16	17	15	22	20	17	22
H.W.	8	10	12	16	18	21	21	18	27	26	22	23
ARI	20	30	54	72	119	115	—	113	160	190	138	142
ALI	22	30	60	69	120	121	100	110	155	202	120	163
ARII	26	33	63	77	119	115	106	134	158	—	—	179
ALII	24	32	59	76	105	—	106	114	156	230	163	—
ARIII	23	30	51	59	97	101	108	108	96	146	121	138
ALIII	23	31	65	—	140	135	—	—	141	229	—	155
ARIV	22	27	54	70	—	123	118	118	157	202	138	162
ALIV	25	30	56	70	152	125	101	115	172	210	142	150
HcAL	23	30	51	59	97	101	108	108	96	146	121	138
Lig. L.	0.8	3.6	4.0	8.3	16.2	17.5	13.3	14.5	17.0	19.0	19.3	20
Cal. L.	0.1	0.7	0.9	2.1	3.6	3.3	2.8	2.7	3.3	3.4	3.1	4.0
Web A	6	8	13	15	19	23	24	17	33	36	21	28
Web BR	7	9	16	17	23	26	27	27	39	41	28	32
BL	7	9	17	17	20	28	23	27	35	47	30	35
Web CR	8	9	18	16	22	29	35	28	37	45	33	36
CL	8	9	17	17	22	33	32	28	38	48	32	37
Web DR	7	10	17	18	22	33	35	30	34	44	37	35
Web DL	7	10	15	16	25	28	27	29	40	45	33	37
Web E	7	8	11	14	20	24	24	25	32	37	27	27
Web Form.	CBDEA	DCBAE	CDBAE	CDBAE	DCBEA	CDBEA	CDBAE	DCBEA	DCBAE	CDBEA	DCBEA	CDBAE
S normal	1.2	1.5	1.9	2.7	3.3	3.6	3.6	4.0	4.5	4.8	4.5	4.8
Sp. L.	—	—	—	—	—	—	36.0	31	—	49	45	42
Sp. R.L.	—	—	—	—	—	—	16.5	14	—	20	13	18
Sp. R.W.	—	—	—	—	—	—	1.6	1.1	—	1.0	1.3	1.3
Penis L.	2	3	3	—	10	—	16	17	14	13	16	17
Gill No.	8	7	7	7	7	8	7	7	8	7	8	7

Sex	♂	♂	♂	♀	♀	♀	♀	♀	♀	♀	♀	♀	♀
M.L.	67	72	73	9	14	17	17	19	22	25	28	34	34
Tot. L.	250	395	285	29	57	57	75	57	87	88	82	121	140
MW	49	65	37	9	12	14	14	14	19	19	26	22	24
H.L.	20	22	24	3	7	6	7	8	7	10	11	13	12
H.W.	29	34	25	7	10	11	10	12	13	15	16	18	16
ARI	172	326	201	15	37	31	—	—	47	56	65	—	86
ALI	175	—	—	16	38	33	52	37	50	59	65	78	79
ARII	187	325	202	18	37	32	35	43	55	62	—	85	—
ALII	206	368	229	19	40	35	43	44	58	—	76	86	109
ARIII	—	205	162	18	39	37	57	40	59	59	65	86	101
ALIII	205	352	215	18	43	36	50	42	58	65	76	76	110
ARIV	—	285	190	18	33	36	53	41	—	61	—	83	101
ALIV	186	300	—	18	39	34	53	—	54	59	73	80	92
HcAL	—	205	162	—	—	—	—	—	—	—	—	—	—
Lig. L.	—	27.6	21.8	—	—	—	—	—	—	—	—	—	—
Cal. L.	—	3.5	2.6	—	—	—	—	—	—	—	—	—	—
Web A	38	55	40	5	10	9	10	10	11	10	15	19	16
Web BR	39	60	42	6	10	10	11	10	15	13	18	22	18
BL	50	61	50	6	12	10	11	10	16	14	18	22	19
Web CR	38	60	44	6	10	12	12	11	17	19	19	25	20
CL	65	60	50	6	12	11	14	12	18	17	20	26	21
Web DR	40	54	46	6	12	10	12	12	17	17	16	24	23
DL	55	55	52	6	11	11	14	13	16	17	20	22	24
Web E	38	43	40	5	11	9	11	9	14	14	16	18	18
Web Form.	CDBAE	BCDAE	DCBAE	DCBAE	DCBEA	CDBAE	DCBEA	DCBAE	CDBEA	CDBEA	CDBEA	CDBAE	DCBEA
S normal	6.7	7.1	5.4	0.7	1.3	1.6	1.4	1.7	2.4	2.3	2.5	2.8	2.3
Sp. L.	—	—	41	—	—	—	—	—	—	—	—	—	—
Sp. R.L.	—	—	16	—	—	—	—	—	—	—	—	—	—
Sp. R.W.	—	—	1.4	—	—	—	—	—	—	—	—	—	—
Penis L.	16	19	19	—	—	—	—	—	—	—	—	—	—
Gill No.	8	7	7	8	7	7	7	7	7	7	8	8	7

Sex	♀	♀	♀	♀	♀	♀	♀	♀	♀	♀
M.L.	37	37	41	44	47	49	49	62	88	—
Tot. L.	152	208	180	162	168	215	250	149	248	—
MW	31	33	24	31	27	37	44	40	32	—
H.L.	11	13	15	14	11	11	14	18	19	—
H.W.	20	21	18	21	18	20	25	23	24	—
ARI	86	136	115	109	100	147	177	120	132	—
ALI	94	140	—	109	—	148	175	122	109	—
ARII	108	159	130	126	119	162	191	141	156	—
ALII	108	156	129	119	113	159	—	133	156	—
ARIII	112	146	131	125	115	162	201	140	166	—
ALIII	106	147	134	117	124	161	198	136	163	—
ARIV	106	132	135	—	113	158	186	126	154	—
ALIV	107	158	—	118	125	157	179	134	162	—
HcAL	—	—	—	—	—	—	—	—	—	—
Lig. L.	—	—	—	—	—	—	—	—	—	—
Cal. L.	—	—	—	—	—	—	—	—	—	—
Web A	20	23	21	26	21	25	35	26	26	—
Web BR	26	31	19	26	28	31	41	29	38	—
BL	27	31	25	30	28	28	35	29	35	—
Web CR	29	35	29	29	30	35	45	28	48	—
CL	27	34	29	29	28	35	47	36	50	—
Web DR	26	33	26	32	27	34	40	29	46	—
DL	25	35	30	25	35	39	46	34	42	—
Web E	23	25	21	23	21	30	31	27	30	—
Web Form.	CBDEA	CDBEA	DCBAE	DCBAE	DCBAE	DCBEA	CDBAE	CDBEA	CDBEA	—
S. normal	2.9	3.4	3.1	3.2	3.1	3.7	4.6	3.8	4.6	—
Sp. L.	—	—	—	—	—	—	—	—	—	—
Sp. R.L.	—	—	—	—	—	—	—	—	—	—
Sp. R.W.	—	—	—	—	—	—	—	—	—	—
Penis L.	—	—	—	—	—	—	—	—	—	—
Gill No.	8	7	7	7	7	7	7	8	7	—

TABLE 2
Means, Standard Deviations and Ranges of Indices of *Octopus australis* Hoyle

	Males				Females			
	n	mean	S.D.(n - 1)	range	n	mean	S.D.(n - 1)	range
MWI	15	73.3	11.8	51-90	19	75.8	15.2	36-100
HLI	15	35.3	6.6	23-48	19	33.5	7.2	22-50
HWI	15	48.8	9.5	34-62	19	53.0	12.4	27-77
ALI	15	78.5	6.9	68-93	19	75.4	7.7	65-95
WDI	15	24.2	4.8	16-32	19	26.8	3.2	22-32
SI (normal)	15	8.9	0.1	7-10	19	8.1	1.3	5-11
HcAI	6	68.2	6.6	58-75				
LLI	11	14.7	1.9	12-18				
CLI	11	18.5	3.9	12-25				
PLI	10	28.5	4.6	23-37				
SpL (mm)	6	40.7	6.4	31-49				
SpLI	6	73.1	12.8	56-91				
SpRI	6	39.3	6.4	29-46				
SpWI	6	3.1	0.6	2.0-3.9				

Explanation of Plate

PLATE 1

Octopus australis Hoyle, 1885. Holotype, female, 22 mm

M.L. a. Dorsal view. b. Lateral view.

a



b

